

SUGARBERRY (CELTIS LAEVIGATA) DIEBACK AND MORTALITY: FINDINGS AND FUTURE WORK

Emilee M. Poole, Michael D. Ulyshen, Scott Horn

US Forest Service, Southern Research Station, Athens, GA

SUGARBERRY



THE PROBLEM



2009: Many dead and dying sugarberry reported in Columbia, SC



High numbers of a buprestid and woolly aphid are found on stressed sugarberry trees and little is known about the insects agent(s)



Causal agent(s) and geographical extent of symptomatic trees are unknown

SYMPTOMS


Small leaves

Thin crowns

Yellowing foliage

Branch dieback

Early leaf senescence







- Large number of egg masses
- Weeping fluid on bark around egg masses
- Staining and galleries with unknown buprestid larvae (later identified)

AGRILUS MACER FLATHEADED HACKBERRY BORER


Historically present in low numbers in southern US- opportunistic

Some dying trees are not attacked by the flatheaded hackberry borer, and others overcome attack

Beetle attacks may hasten the death of the trees, and prevent recovery

A. macer is not the primary cause of mortality









NON-NATIVE APHIDS: SHIVAPHIS CELTI IN THE NEWS

What are those snowflakes falling out of the sky?

September Snow? Woolly aphids here until end of summer

It's not snowing: those white things are woolly aphids!

An aphid snowstorm covers area again

Early 'snow' proves to be woolly aphids



- Chlorosis following feeding
- Honeydew & sooty mold
- Premature leaf drop
- Repeated yearly damage

SHIVAPHIS CELTI: HACKBERRY WOOLLY APHID

Imidacloprid consistently provides protection against hackberry woolly aphids throughout the growing season for sugarberry

In the US, most hackberry woolly aphid reports are in the southeast. Aphid is likely present in areas with *Celtis*

Impact of hackberry woolly aphid is still poorly understood



MORTALITY
CONTINUES TO
EXPAND IN THE
SOUTHEASTERN
US




OBJECTIVES

Document patterns of dieback and mortality over five years of observation

Determine the efficacy of systemic insecticide treatments in effort to improve survivorship

Report known locations with high mortality



Symptomatic: 3 or higher




- Stem injection of emamectin benzoate on October 22, 2015 caused bark damage.
- Imidacloprid soil injections applied:

October 22, 2015, April 3, 2017, and April 22, 2020


• Mortality did not differ among the insecticide treatment groups($X^2[3, n = 59] = 1.02, p = 0.80$) based on logistic regression


RESULTS AFTER 5 YEARS OF MONITORING

~52% of monitored trees died in 5 years





SURVIVAL TIME AFTER BECOMING SYMPTOMATIC






CITIES WITH HIGH MORTALITY









FINAL CONCLUSIONS

Compared to other decline/dieback events this is one of the most long-lasting and expansive *Celtis* mortality events

No signs of the dieback and decline showing; primary cause still unknown



- Canopy sampling
- Celtis determination
- Phenolic data analysis
- Reassess monitoring sites
- Native/invasive understory recruitment
- Local climate role



NEW OPPORTUNITIES AND FUTURE WORK





SUGARBERRY SQUAD IN ACTION



























Ulyshen tag height

Emilee

ACKNOWLEDGMENTS

- Roy Kibler (North Augusta Superintendent of Property Maintenance)
- Lee Herman (American Museum of Natural History)
- Melissa Callahan (Auburn University Natural History Museum)
- Jacqueline Airoso (California State Collection of Arthropods)
- Patrice Bouchard and Anthony Davies (Canadian National Collection of Insects, Arachnids, and Nematodes)
- John Rawlings and Robert Androw (Carnegie Museum of Natural History Invertebrate Collection)
- John Morse and Mike Ferro (Clemson University Arthropod Collection)
- Christopher Grinter (Entomology Collection of the California Academy of Sciences)
- Jim Louderman and Crystal Maier (Field Museum of Natural History Collection of Insects, Arachnids, and Myriapods and InvertEbase)
- Rick Hoebeke (Georgia Natural History Museum)
- Philip Perkins (Harvard Museum of Comparative Zoology)
- Tommy McElrath (Illinois Natural History Survey Insect Collection)
- Louisiana State Arthropod Museum
- Gary Parsons and Anthony Cognato (Albert J. Cook Arthropod Research Collection at Michigan State)
- Terence Lee Schiefer (Mississippi Entomological Museum)

Weiping Xie and Brian Brown (Natural History Museum of Los Angeles County)

- Hellen Vessels (New Mexico State University Arthropod Collection)
- Bob Blinn (North Carolina State University Insect Collection)
- Luciana Musetti (Ohio State Triplehorn Insect Collection)
- Phil Mulder and Jana Slaughter (Oklahoma State- K.C. Emerson Entomology Museum)
- Charyn J. Micheli and Eugenio H. Nearns (Smithsonian National Museum of Natural History)
- Karen Wright (Texas A&M Insect Collection)
- Texas Tech Insect Collection
- Wendy Moore and Wesley E. Hall (University of Arizona)
- Michael S. Engel (University of Kansas Natural History Museum)
- Stylianos Chatzimanolis (University of Tennessee Natural History Museum)
- Stephen Fraedrich, Michelle Cram, and Rabiu Otalinwo (USDA FS)
- Dave Coyle (Clemson University)
- Chris Asaro, Chip Bates, Chris Barnes, Lynn Womack, Alex Ballard, Seth Hawkins (GFC), David Jenkins (SFC)
- David Shelley (National Park Service)
- Villari, Batzer, Aubrey, Ulyshen, and Braman Labs

USDA Forest Service, Southern Research Station and Forest Health Protection's Evaluation Monitoring program (SO-EM-17-04)



Filmed by M. D. Ulyshen

Thank you Emilee.poole@usda.gov